

A. Mature Tm 13.17 amino acid residue

1 LTEAQIEKLN KISKKCQNES GVSQEIITKA RNGDWEDDPK LKRQVFCVAR
51 NAGLATESGE VVVDVLREKV RKVTDNDEET EKIINKCAVK RDTVEETVFN
101 TFKCVMKNKP KFSPVD

B. Summary of the composition analysis for the mature Tm 13.17 sequence:

<u>Residue</u>	<u>Number</u>	<u>Mole Percent</u>
A = Ala	6	5.172
B = Asx	0	0.000
C = Cys	4	3.448
D = Asp	8	6.897
E = Glu	13	11.207
F = Phe	4	3.448
G = Gly	4	3.448
H = His	0	0.000
I = Ile	6	5.172
K =Lys	16	13.793
L = Leu	5	4.310
M = Met	1	0.862
N = Asn	8	6.897
P = Pro	3	2.586
Q = Gln	4	3.448
R = Arg	6	5.172
S = Ser	5	4.310
T = Thr	8	6.897
V =Val	14	12.069
W = Trp	1	0.862
Y = Tyr	0	0.000
Z = Glx	0	0.000

Molecular weight = 13171.96; Residues = 116; Average Residue Weight = 113.551

Charge = 1; Isoelectric point = 7.74.

FIG 2.6c